





# Responsible RestartOhio

# **Manufacturing, Distribution & Construction**







# **Employees**, Distributors. & Guests

## **Mandatory**

- Ensure minimum 6 ft between people, if not possible, install barriers
- Employees must perform daily symptom assessment\*
- Require employees to stay home if symptomatic
- Face coverings must be worn at all times
- Require regular handwashing
- · Stagger or limit arrivals of employees and guests
- Personnel should work from home if possible

#### **Recommended Best Practices**

 Provide stipend to employees for private transportation

#### **Shift Pattern**

- Daily disinfection of desks and workstations
- Change shift patterns (e.g. fewer shifts)
- Stagger lunch and break times

- Split into sub-teams, limit contact across sub-teams
- Reduce pace to allow less FTEs per line

# **Physical** Spaces / Workstations

- Ensure minimum 6 ft between people, if not possible, install barriers
- Daily deep disinfection of high-contact surfaces
- · Space factory floor to allow for distancing
- · Regulate max number of people in cafeterias/ common spaces
- · Establish maximum capacity (e.g. 50% of
- · Close cafeteria and gathering spaces if possible, or conduct regular cleanings
- · Daily deep disinfection of entire facility

### Confirmed Cases

- Immediately isolate and seek medical care for any individual who develops symptoms while at work
- · Contact the local health district about suspected cases or exposures
- Shutdown shop/floor for deep sanitation if possible
- · Work with local health department to identify potentially infected or exposed individuals to help facilitate effective contact tracing/notifications
- Once testing is readily available, test all suspected infections or exposures
- Following testing, contact local health department to initiate appropriate care and tracing







# Responsible RestartOhio

# **Consumer, Retail & Services**







## **Employees**

## Mandatory

- Ensure min 6 ft between people, if not possible, install barriers
- Employees must perform daily symptom assessment\*
- · Require employees to stay home if symptomatic
- Face coverings must be worn at all times
- Require regular handwashing
- Place hand sanitizers in high-contact locations
- Clean high-touch items after each use (e.g. carts, baskets)

#### Recommended Best Practices

Group employees by shift to reduce exposure

### Customers & Guests

#### Ensure minimum 6 ft between people

- Specify hours for at-risk populations (e.g. elderly)
- Place hand sanitizers in high-contact locations
- Ask customers and guests not to enter if symptomatic
- Face coverings to be worn while shopping or visiting
- Stagger entry of customers and guest

- Health questionnaire for symptoms at entry point
- Provide face coverings upon entry
- Where possible, accept customers by appointment only
- Increase availability for curb-side pickup
- Consider suspending return policies

# **Physical Spaces**

#### Ensure minimum 6 ft between people, if not possible, install barriers

- Post social distancing signage & disinfect high-contact surfaces hourly
- Clean merchandise before stocking if possible
- Establish maximum capacity (e.g. 50% of fire code)
- Discontinue self-service food stations, product samples
- Food courts remain closed

- Close once a week for deep cleaning
- Maximize available checkout space to promote social distancing (e.g., space customer lines with floor markers, use alternate registers)
- Use contact-less payments where possible
- Increase capacity for delivery and curb-side pickup

## Confirmed Cases

- Immediately isolate and seek medical care for any individual who develops symptoms while at work
- Contact the local health district about suspected cases or exposures
- · Shutdown shop/floor for deep sanitation if possible
- Work with local health department to identify potentially infected or exposed individuals to help facilitate effective contact tracing/ notifications
- Once testing is readily available, test all suspected infections or exposures
- Following testing, contact local health department to initiate appropriate care and tracing

<sup>\*</sup>Daily symptom assessments should include taking your temperature with a thermometer and monitoring for fever. Also watch for coughing or trouble breathing.







# Responsible RestartOhio

# **General Office Environments**







#### **Mandatory**

# **Employees &** Guests

- Ensure minimum 6 ft between people, if not possible, install barriers
- Personnel should work from home when possible
- Employees must perform daily symptom assessment\*
- Require employees to stay home if symptomatic
- Face coverings must be worn at all times while working
- Require regular handwashing
- Reduce sharing of work materials
- Limit travel as much as possible
- Stagger arrival of all employees and guests
- · Post signage on health safety guidelines in common areas

# Recommended Best Practices

- Ensure seating distance of 6 ft or more
- Enable natural workplace ventilation
- Health questionnaire for symptoms at entry
- Temperature taking protocol

# **Physical** Spaces / Workstations

- Frequent disinfection of desks, workstations, and high-contact surfaces
- Daily disinfection of common areas
- Cancel/postpone in person events when social distancing guidelines cannot be met
- No buffet in cafeteria
- Utilize disposable tableware and other materials
- Establish maximum capacity (e.g. 50% of fire code)

#### • Redesign/space workstations for 6 ft or more of distance

- · Close cafeteria and gathering spaces if possible, or conduct regular cleanings
- Limit congregation in office spaces
- Divide essential staff into groups and establishing rotating shift
- Availability of at least 3 weeks of cleaning supplies

# **Confirmed** Cases

- Immediately isolate and seek medical care for any individual who develops symptoms while at work
- Contact the local health district about suspected cases or exposures
- Shutdown shop/floor for deep sanitation if possible
- Work with local health department to identify potentially infected or exposed individuals to help facilitate effective contact tracing/notifications
- Once testing is readily available, test all suspected infections or exposures
- Following testing, contact local health department to initiate appropriate care and tracing